




A Fixed-Point Algorithm for Computing Robust Barycentres of Measures

Eloi Tanguy, Julie Delon, Nathaël Gozlan

1.1 The Monge Problem




$$\begin{aligned}\mu &:= \frac{1}{n} \sum_{i=1}^n \delta_{x_i}, \quad \nu := \frac{1}{n} \sum_{i=1}^n \delta_{y_i} \\ \mathcal{T}_c(\mu, \nu) &:= \min_{\sigma \in \mathfrak{S}_n} \frac{1}{n} \sum_{i=1}^n c(x_i, y_{\sigma(i)}). \\ \text{▪ } p, q \text{ cost: } c_{p,q}(x, y) &:= \|x - y\|_p^q. \\ \text{▪ } q\text{-Wasserstein: } W_q^q &:= \mathcal{T}_{p=2, q=q}.\end{aligned}$$

1.2 2-Wasserstein Barycentres

Euclidean Fréchet mean: $\operatorname{argmin}_{x \in \mathbb{R}^d} \sum_{k=1}^K \lambda_k \|x - y_k\|_2^2 = \sum_{k=1}^K \lambda_k y_k$.

W_2^2 barycentre (Aguech and Carlier 2011): $\operatorname{argmin}_{\mu \in \mathcal{P}(\mathbb{R}^d)} \sum_{k=1}^K \lambda_k W_2^2(\mu, \nu_k)$.



1.3 Fixed-Point Method for W_2^2 barycentres

- Discrete case: Cuturi and Doucet 2014.
- Proofs for absolutely continuous case: Álvarez-Esteban et al. 2016.

Objective: $\operatorname{argmin}_{(x_i)_{i=1}^n} \sum_{k=1}^K \lambda_k \min_{\sigma_k \in \mathfrak{S}_n} \sum_{i=1}^n \|x_i - y_{k, \sigma_k(i)}\|_2^2 = \sum_{k=1}^K \lambda_k W_2^2(\mu, \nu_k)$.

Fixed-Point Algorithm for W_2^2 Barycentres

Input: For $k \in [1, K]$, target $(y_{k,i})_{i=1}^n$. Weights (λ_k) .

```

1 for t ∈ [0, T − 1] do
2   For k ∈ [1, K], σk ∈ argminσ ∈ Sn ∑i=1n ‖xi(t) − yk,σ(i)‖22;
3   For i ∈ [1, n], xi(t+1) = ∑k=1K λkyk,σk(i);
```

2.1 Barycentres for Generic Transport Costs

Objective: $\operatorname{argmin}_{(x_i)_{i=1}^n} \sum_{k=1}^K \lambda_k \min_{\sigma_k \in \mathfrak{S}_n} \sum_{i=1}^n c_k(x_i, y_{k, \sigma_k(i)}) = \sum_{k=1}^K \lambda_k \mathcal{T}_{c_k}(\mu, \nu_k)$.

Fixed-Point Algorithm for OT Barycentres

Input: For $k \in [1, K]$, target $(y_{k,i})_{i=1}^n$. Weights (λ_k) .

```

1 for t ∈ [0, T − 1] do
2   For k ∈ [1, K], σk ∈ argminσ ∈ Sn ∑i=1n ck(xi(t), yk,σ(i));
3   For i ∈ [1, n], xi(t+1) = argminx ∑k=1K λkck(x, yk,σk(i));
```

Long paper: Tanguy et al. 2024 for formulation with generic measures.

2.2 Algorithm Convergence

- Assuming only continuous costs and measures on compact metric spaces.
- Extends Álvarez-Esteban et al. 2016.

Energy function: $V(\mu) := \sum_{k=1}^K \lambda_k W_2^2(\mu, \nu_k)$. Iterates: $\mu_{t+1} \in G(\mu_t)$.

Decrease Property

$$\forall \bar{\mu} \in G(\mu), V(\mu) \geq V(\bar{\mu}) + \mathcal{T}_\delta(\mu, \bar{\mu}).$$


If μ^* is a barycentre then $G(\mu^*) = \{\mu^*\}$.

Convergence

If μ is a subsequential limit of (μ_t) then $\mu \in G(\mu)$.

3.1 3D Barycentre of 2D Point Clouds

$$\text{Objective: } \operatorname{argmin}_{(x_i)_{i=1}^n \in \mathbb{R}^{n \times 3}} \frac{1}{K} \sum_{k=1}^K \min_{\sigma_k \in \mathfrak{S}_n} \sum_{i=1}^n \|P_k x_i - y_{k, \sigma_k(i)}\|_2.$$





3.2 Robust Barycentres


$$c(x, y) := \|x - y\|_{1.5}^{1.5}$$

1.5-Wasserstein barycentre,

Robust to (heart) outlier measure.



3.3 Colour Transfer With Outliers



- [1] Martial Aguech and Guillaume Carlier. "Barycenters in the Wasserstein Space". In: *SIAM Journal on Mathematical Analysis* 43.2 (2011), pp. 904–924.
- [2] Pedro C Álvarez-Esteban, E Del Barrio, JA Cuesta-Albertos, and C Matrán. "A fixed-point approach to barycenters in Wasserstein space". In: *Journal of Mathematical Analysis and Applications* 441.2 (2016), pp. 744–762.
- [3] Marco Cuturi and Arnaud Doucet. "Fast Computation of Wasserstein Barycenters". In: *Proceedings of the 31st International Conference on Machine Learning*. Vol. 32. Proceedings of Machine Learning Research. Beijing, China: PMLR, June 2014, pp. 685–693.
- [4] Eloi Tanguy, Julie Delon, and Nathaël Gozlan. *Computing Barycentres of Measures for Generic Transport Costs*. 2024. arXiv: 2501.04016 [math.NA].